

Last updated: 21 April, 2015

## LAPHYRAGOGUS

*Laphyragogus* Kohl, 1889b:190. Type species: *Laphyragogus pictus* Kohl, by monotypy.

*Leianthrena* Bingham in Kohl, 1897a:381. Type species: *Leianthrena kohlii* Bingham, 1897, by original designation.

*Leianthrena* Bingham, 1897:212. Type species: *Leianthrena kohlii* Bingham, by original designation [= *Leianthrena kohlii* Bingham in Kohl, 1897].

Revision: de Beaumont, 1959d.

Review of biology: Kazenas, 2001b:146.

### 1. *ajjer* de Beaumont

*Laphyragogus ajjer* de Beaumont, 1958b:63, ♂. Holotype: ♂, Algeria: Tassili des Ajjer: reg between oueds Sersouf and Issandilène (MNHN). – de Beaumont, 1959d:732 (in revision of *Laphyragogus*); R. Bohart and Menke, 1976:220 (in checklist of world Sphecidae); Prentice, 1993:7 (possibly preying on adult Lepidoptera).

### *ssp. orientalis* de Beaumont

*Laphyragogus ajjer orientalis* de Beaumont, 1970c:17, ♂, ♀. Holotype: ♂, Iran: Baluchistan: SW Iranshar (Ludwigsburg). – Bohart and Menke, 1976:220 (in checklist of world Sphecidae).

### 2. *kohlii* (Bingham)

*Leianthrena kohlii* Bingham in Kohl, 1897:381, ♀, ♂ (as *Kohlii*, incorrect original capitalization). Syntypes: Pakistan: North-West Provinces: no specific locality, and India or Pakistan: Punjab: no specific locality (BMNH – **As *Laphyragogus kohlii***: de Beaumont, 1959d:733 (in revision of *Laphyragogus*); R. Bohart and Menke, 1976:220 (in checklist of world Sphecidae); Kazenas, 2001:25 (in checklist of Sphecidae of Kazakhstan and Central Asia).

*Lianthrena kohlii* Bingham, 1897:213, ♀, ♂. Syntypes: India or Pakistan: Punjab: no specific locality (BMNH). Objective synonym of *Leianthrena kohlii* Bingham, 1897.

*Laphyragogus turanicus* Gussakovskij, 1952:227, ♀, ♂. Lectotype: ♀, Tajikistan: Ayvadj at Kafirnigan River (ZIN), designated by Kazenas, 2000a:50. Synonymized with *Laphyragogus kohlii* by Kazenas, 2000a:50. – de Beaumont, 1959d:733 (in revision of *Laphyragogus*); R. Bohart and Menke, 1976:220 (in checklist of world Sphecidae); Kazenas, 1985:17 (Tajikistan; prey: adult *Crambus*), Kazenas, 2001b:146 (review of known biology).

### 3. *pectinatus* de Beaumont

*Laphyragogus pectinatus* de Beaumont, 1959d:728, ♀, ♂. Holotype: ♂, Israel: Revivim (Lausanne). – de Beaumont, 1960b:246 (Libya); de Beaumont, Bytinski-Salz and Pulawski, 1973:18 (Israel); R. Bohart and Menke, 1976:220 (in checklist of world Sphecidae); Roche, 2007a:61 (in checklist of Egyptian Crabronidae, redescription), 2007b:4 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae).

### 4. *pictus* Kohl

*Laphyragogus pictus* Kohl, 1889:190, ♀. Holotype: Egypt: no specific locality (ZMHU). – Mochi, 1938:223 (Egypt; description of ♂); Giner Marí, 1945e:235 (Spanish Sahara); de Beaumont, 1959d:724 (in revision of western Palearctic *Laphyragogus*); de Beaumont, Bytinski-Salz and Pulawski, 1973:18 (Israel); R. Bohart and Menke, 1976:220 (in checklist of world Sphecidae); Radović, 1985:64 (sting apparatus analyzed); Al-Houty, 1989:160 (Kuwait: Al-Wafra); Hua, 2006:280 (in list of Chinese insects, geographic distribution); Roche, 2007a:62 (in checklist of Egyptian Crabronidae), 2007b:4 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae).

**5. *strakai* Schmid-Egger**

*Laphyragogus strakai* Schmid-Egger, 2011b:543, ♂. Holotype: ♂, United Arab Emirates: Um al-Quwain at 25°32'N 55°32'E (Ch. Schmid-Egger personal collection, Berlin). – Schmid-Egger, 2011b:544 (color photograph of male).

**6. *visnagae* de Beaumont**

*Laphyragogus visnagae* de Beaumont, 1959d:730, ♀, ♂. Holotype: ♂, Algeria: Biskra (Lausanne). – Bohart and Menke, 1976:220 (in checklist of world Sphecidae).

**sp.**

de Beaumont, 1957a:143 (Morocco), 1961b:275 (Afghanistan); Roche, 1981:2 (in checklist of Sphecidae of United Arab Emirates); Schmid-Egger, 2011b:545 (United Arab Emirates, description of ♀).